


ABCD liraglutide audit HbA1c changes according to duration of diabetes

Figure 2. Mean HbA_{1c} changes after 26 weeks of liraglutide treatment, stratified according to duration of diabetes




Columns show adjusted mean changes analysed by ANCOVA with baseline HbA_{1c} as a covariate. ED: estimated difference; CI: confidence interval

Estimated change in HbA_{1c} and body weight by baseline diabetes duration

Diabetes duration

Change from baseline:

	≤5 years				>5 to ≤10 years				>10 years			
	HbA _{1c} (%)		Body weight (kg)		HbA _{1c} (%)		Body weight (kg)		HbA _{1c} (%)		Body weight (kg)	
N	533	641	533	641	423	565	423	565	376	528	376	528
Baseline	8.1	8.1	95.9	95.9	8.2	8.2	93.5	93.5	8.3	8.3	89.8	89.8
End of treatment	6.6	6.4	91.8	90.1	6.8	6.4	89.6	87.7	6.8	6.5	85.8	84.1



Data presented are estimated change from baseline to week 30 or week 40 based on a meta-analysis of data from the six trials. BW, body weight; N, number of subjects in the full analysis set. Rosenstock J et al. Presented at the 78th Scientific Sessions of the American Diabetes Association, 22–26 June, 2018, Orlando, Florida, USA: Poster Presentation 1081-P.