


QUICK REFERENCE GUIDE FOR SURGICAL DIABETES MANAGEMENT

A CHART MUST BE FOLLOWED ONCE PATIENT HAS MISSED ONE MEAL



NOTE FOR FLUIDS IN VRIII *

For patients with hip fractures, CKD fluid restriction or a history/signs of heart failure:

Standard fluid should be given at 50ml/hr
(**not** 100ml/hr)

There are two ways to manage peri-operative starvation for patients with diabetes

Drug Management (Drug Mx) =

- Not on VRIII (Sliding scale)
- Always continue Long acting insulin as explained on Drug Management chart
- Only continue Short acting / Pre-mixed insulin with Pre-Assessment Clinic (PAC) diabetes plan

Patient's own Insulin Pump =

- Continuous Subcutaneous Insulin Infusion = CSII
- Look for plan on CWS, discuss with patient
- Start **Drug Management** chart
- Continue basal
- Stop bolus
- See intranet guidance

Variable Rate IV Insulin Infusion (VRIII) =

- Previously known as Sliding scale
- Always continue long acting insulin as per VRIII chart instructions (page 1 of chart)

IF IN DOUBT CALL FOR HELP

Diabetes nurses

Richmond House

09:00 -17:00 ext 44271

If no answer leave message

In an emergency call:

Anaes Reg

Med Reg